arXiv:1102.0662v4 [math.NA] 17 Dec 2011

The tamed Milstein method for commutative stochastic differential equations with non-globally Lipschitz continuous coefficients *

Siqing Gan Xiaojie Wang †

School of Mathematical Sciences and Computing Technology, Central South University,
Changsha 410075, Hunan, PR China

Abstract

For stochastic differential equations (SDEs) with a superlinearly growing and globally one-sided Lipschitz continuous drift coefficient, the classical explicit Euler scheme fails to converge strongly to the exact solution. Recently, an explicit strongly convergent numerical scheme, called the tamed Euler method, is proposed in [Hutzenthaler, Jentzen, & Kloeden (2010); Strong convergence of an explicit numerical method for SDEs with non-globally Lipschitz continuous coefficients, arXiv:1010.3756v1] for such SDEs. Motivated by their work, we here introduce a tamed version of the Milstein scheme for SDEs with commutative noise. The proposed method is also explicit and easily implementable, but achieves higher strong convergence order than the tamed Euler method does. In recovering the strong convergence order one of the new method, new difficulties arise and kind of a bootstrap argument is developed to overcome them. Finally, an illustrative example confirms the computational efficiency of the tamed Milstein method compared to the tamed Euler method.

AMS subject classification: 65C20, 60H35, 65L20.

Key Words: tamed Milstein method, superlinearly growing coefficient, one-sided Lipschitz condition, commutative noise, strong convergence

1 Introduction

We consider numerical integration of stochastic differential equations (SDEs) in the Itô's sense

$$dX_t = \mu(X_t)dt + \sigma(X_t)dW_t, \quad X_0 = \xi, \quad t \in [0, T]. \tag{1.1}$$

---

[*] This work was supported by NSF of China (No.11171352) and Hunan Provincial Innovation Foundation For Postgraduate (NO.CX2010B118)

[†] Corresponding author: x.j.wang7@gmail.com

Here $\mu : \mathbb{R}^d \longrightarrow \mathbb{R}^d, \sigma = (\sigma_1, \sigma_2, ..., \sigma_m) : \mathbb{R}^d \longrightarrow \mathbb{R}^{d\times m}$. We assume that W_t is an m-dimensional Wiener process defined on the complete probability space $(\Omega, \mathcal{F}, \mathbb{P})$ with an increasing filtration $\{\mathcal{F}_t\}_{t\geq 0}$ satisfying the usual conditions. And the initial data ξ is independent of the Wiener process. (1.1) can be interpreted mathematically as a stochastic integral equation

$$X_t = X_0 + \int_0^t \mu(X_s)ds + \sum_{i=1}^{m} \int_0^t \sigma_i(X_s)dW_s^i, \quad t \in [0, T], \mathbb{P} - a.s., \tag{1.2}$$

where $\sigma_i(x) = (\sigma_{1,i}(x), ..., \sigma_{d,i}(x))^T$ for $x \in \mathbb{R}^d, i \in \{1, 2, ..., m\}$ and the second integral is the Itô integral.

This article is concerned with the strong approximation problem (see, e.g., Section 9.3 in Kloeden and Platen [12]) of the SDEs (1.2). More precisely, on a uniform mesh with stepsize $h = \frac{T}{N}$ defined by $\mathcal{T}^N : 0 = t_0 < t_1 < t_2 < \cdots < t_N = T, N \in \mathbb{N}$, we want to compute a numerical approximation $Y_n : \Omega \to \mathbb{R}^d, n \in \{0, 1, ..., N\}$ with $Y_0 = \xi$ such that

$$\left(\mathbb{E}\|X_T - Y_N\|^2\right)^{\frac{1}{2}} < \varepsilon \tag{1.3}$$

for a given precision $\varepsilon > 0$ with the least possible computational effort. The strong convergence problem becomes very important because efficient Multi-Level Monte Carlo (MLMC) simulations rely on the strong convergence properties [3].

The simplest and most obvious idea to solve the strong approximation problem (1.3) is to apply the explicit Euler scheme [15]

$$Y_{n+1} = Y_n + h\mu(Y_n) + \sigma(Y_n)\Delta W_n, \quad Y_0 = \xi,\ n = 0, 1, ..., N - 1, \tag{1.4}$$

where $\Delta W_n = W_{t_{n+1}} - W_{t_n}$. The method is strongly convergent with order one half if the coefficients μ, σ satisfy the global Lipschitz condition (see, for instance, [12]). Unfortunately, it has recently been shown in [7] that the explicit Euler scheme fails to provide strong convergent solution to the SDEs with super-linearly growing drift coefficient. It is well-known that the backward Euler method can promise strong convergence in this situation, see e.g.,[4]. But the backward Euler method is an implicit method, which requires additional computational effort to solve an implicit system. Recently in [9], the authors proposed an explicit method, called tamed Euler method, for (1.2)

$$Y_{n+1} =Y_n + h\tilde{\mu}(Y_n) + \sigma(Y_n)\Delta W_n, \quad \text{with} \quad \tilde{\mu}(Y_n) = \frac{h\mu(Y_n)}{1 + h\|\mu(Y_n)\|}. \tag{1.5}$$

Here $\tilde{\mu}$ is a modification of μ. This tamed Euler scheme is proved to converge strongly with the standard convergence order 0.5 to the exact solution of (1.2) if the drift coefficient function is globally one-sided Lipschitz continuous and has an at most polynomially growing derivative.

On the one hand, the explicit Milstein scheme is another numerical scheme for SDEs that achieves a strong order of convergence higher than that of the explicit Euler scheme (1.4). In fact the explicit Milstein scheme has strong convergence order of one if the coefficient functions of stochastic Taylor expansions satisfy both the global Lipschitz condition and the linear growth condition(see [12]). The explicit Milstein method [12, 16] applied to (1.1) reads

$$Y_{n+1} =Y_n + h\mu(Y_n) + \sigma(Y_n)\Delta W_n + \sum_{j_1,j_2=1}^{m} L^{j_1}\sigma_{j_2}(Y_n)I_{j_1,j_2}^{t_n,t_{n+1}}, \tag{1.6}$$

where

$$L^{j_1} = \sum_{k=1}^{d} \sigma_{k,j_1} \frac{\partial}{\partial x^k}, \quad I_{j_1,j_2}^{t_n,t_{n+1}} = \int_{t_n}^{t_{n+1}} \int_{t_n}^{s_2} dW_{s_1}^{j_1} dW_{s_2}^{j_2}. \tag{1.7}$$

Since the explicit Milstein scheme and the explicit Euler scheme coincide when applied to the SDEs with additive noise, we can deduce from the results in [7] that the explicit Milstein scheme generally does not converge in the mean-square sense to the exact solution solution of the SDEs with super-linearly growing drift coefficient. Accordingly, we follow the idea from [9] and replace $\mu(Y_n)$ in (1.6) with $\tilde{\mu}(Y_n)$ to derive a tamed Milstein method

$$Y_{n+1} = Y_n + h\tilde{\mu}(Y_n) + \sigma(Y_n)\Delta W_n + \sum_{j_1,j_2=1}^{m} L^{j_1}\sigma_{j_2}(Y_n) I_{j_1,j_2}^{t_n,t_{n+1}}, \tag{1.8}$$

which we expect to be strongly convergent with order one in the non-globally Lipschitz case.

On the other hand, although Milstein-type schemes may achieve a strong convergence order higher than that of Euler type schemes, additional computational effort is required to approximate the iterated Itô integrals $I_{j_1,j_2}^{t_n,t_{n+1}}$ for every time step [13]. This will enable the Milstein-type schemes to lose their advantage over the Euler-type schemes in computational efficiency. In this article we restrict our attention to SDEs with commutative noise, in which case the Milstein scheme can be easily implemented without simulating the iterated Itô integrals. In this situation, Milstein-type method is much more computational efficient than Euler-type method. More precisely, let the diffusion matrix σ fulfill the so-called commutativity condition:

$$L^{j_1}\sigma_{k,j_2} = L^{j_2}\sigma_{k,j_1}, \quad j_1, j_2 = 1, ..., m, k = 1, ..., d. \tag{1.9}$$

In many applications the considered SDE systems possess commutative noise (see [12]).

Thanks to the property $I_{j_1,j_2}^{t_n,t_{n+1}} + I_{j_2,j_1}^{t_n,t_{n+1}} = \Delta W_n^{j_1} \Delta W_n^{j_2}, j_1 \neq j_2$, in this case the tamed Milstein method (1.8) takes a simple form as

$$Y_{n+1} = Y_n + h\tilde{\mu}(Y_n) + \sigma(Y_n)\Delta W_n + \frac{1}{2}\sum_{j_1,j_2=1}^{m} L^{j_1}\sigma_{j_2}(Y_n)\left(\Delta W_n^{j_1}\Delta W_n^{j_2} - \delta_{j_1,j_2}h\right), \tag{1.10}$$

where $\delta_{j_1,j_2} = 1$ for $j_1 = j_2$ and $\delta_{j_1,j_2} = 0$ for $j_1 \neq j_2$, $\tilde{\mu}$ is the modification of μ as defined in (1.5).

The main result of this article shows that the tamed Milstein scheme (1.10) converges strongly with the standard convergence order one to the exact solution of SDEs with commutative noise if the drift coefficient μ is globally one-sided Lipschitz continuous and has at most polynomially growing first and second derivatives. The diffusion coefficient σ and the coefficient function of stochastic Taylor expansion $L^{j_1}\sigma_{j_2}, j_1, j_2 \in \{1, ..., m\}$ are assumed to be globally Lipschitz continuous. It is worthwhile to mention that a similar approach as used in [9] is evoked to obtain uniform boundedness of p-th moments of numerical solutions produced by the tamed Milstein method. We also introduce similar stochastic processes D_n that dominate the tamed Milstein approximation on appropriate subevents Ω_n (see Section 2 for more details). With bounded p-th moments at hand, our main effort is to show for the time continuous tamed Milstein methods there exists a family of real numbers $C_{p,T} \in [1, \infty)$ for $p \in [1, \infty)$ such that

$$\left(\mathbb{E}\Big[\sup_{t\in[0,T]} \Big\|X_t - \bar{Y}_t\Big\|^p\Big]\right)^{1/p} \leq C_{p,T} \cdot h, \quad h \in (0, 1]. \tag{1.11}$$

The key difficulty is that relative to previous analysis [9] a sharper estimate of the term J (see (3.35)) must be obtained to get the strong convergence order one. To overcome this difficulty, a certain kind of bootstrap argument is exploited (see the estimate of J for more details). To the best of our knowledge, this is the very first paper to successfully recover the strong convergence order one for the Milstein-type method under non-globally Lipschitz condition.

The rest of this paper is arranged as follows. In the next section, uniform boundedness of p-th moments are obtained. And then the strong convergence order of the tamed Milstein method is established in Section 3. Finally, an illustrative example confirms the strong convergence order of one and the computational efficiency of this scheme compared to the tamed Euler scheme.

2 Uniform boundedness of p-th moments

Throughout this article, $N \in \mathbb{N}$ is the step number of the uniform mesh defined in the previous section. Moreover, we use the notation $\|x\| := (|x_1|^2 + ... + |x_k|^2)^{\frac{1}{2}}$, $\langle x, y\rangle := x_1y_1 + ... + x_ky_k$ for all $x = (x_1, x_2, ..., x_k), y = (y_1, y_2, ..., y_k) \in \mathbb{R}^k, k \in \mathbb{N}$, and $\|A\| := \sup_{x\in\mathbb{R}^l,\|x\|\leq 1} \|Ax\|$ for all $A \in \mathbb{R}^{k\times l}, k, l \in \mathbb{N}$. Furthermore, we make the following assumptions.

Assumption 2.1 *Let $\mu(x)$ and $\sigma_i(x), i = 1, ..., m$ be continuously differentiable and there exist positive constants $K \geq 1$ and $c \geq 1$, such that $\forall x, y \in \mathbb{R}^d$*

$$\begin{aligned} \langle x - y, \mu(x) - \mu(y)\rangle &\leq K\|x - y\|^2, && (2.1)\\ \|\sigma(x) - \sigma(y)\| &\leq K\|x - y\|, && (2.2)\\ \|L^{j_1}\sigma_{j_2}(x) - L^{j_1}\sigma_{j_2}(y)\| &\leq K\|x - y\|, \quad j_1, j_2 \in \{1, 2, ..., m\}, && (2.3)\\ \|\mu'(x)\| &\leq K(1 + \|x\|^c). && (2.4) \end{aligned}$$

Note that the globally one-sided Lipschitz condition (2.1) on the drift μ and the globally Lipschitz condition (2.2) on the diffusion σ have been widely used in the literatures, e.g., [4, 5, 6, 7, 8, 9].

To prove uniform boundedness of p-th moments of the numerical solution, we follow the ideas in [9] to introduce the appropriate subevents Ω_n and dominating stochastic processes D_n

$$\Omega_n := \Big\{\omega \in \Omega | \sup_{0\leq k\leq n-1} D_k(\omega) \leq \frac{1}{N^{2c}}, \sup_{0\leq k\leq n-1} \|\Delta W_k\| \leq 1\Big\}, \tag{2.5}$$

$$D_n := (\lambda + \|\xi\|) \exp\Big(\lambda + \sup_{0\leq u\leq n} \sum_{k=u}^{n-1} \big[\lambda\|\Delta W_k\|^2 + \alpha_k\big]\Big), \tag{2.6}$$

where

$$\lambda = \Big(1 + 4TK + 2T\|\mu(0)\| + 2K + 2\|\sigma(0)\| + m^2(T + 1)\Big(K + \max_{1\leq j_1,j_2\leq m} \|L^{j_1}\sigma_{j_2}(0)\|\Big)\Big)^2$$

and

$$\alpha_n = 1_{\{\|Y_n\|\geq 1\}}\Big\langle \frac{Y_n}{\|Y_n\|}, \frac{\sigma(Y_n)}{\|Y_n\|}\Delta W_n\Big\rangle + 1_{\{\|Y_n\|\geq 1\}}\Big\langle \frac{Y_n}{\|Y_n\|}, \sum_{j_1,j_2=1}^{m} \frac{L^{j_1}\sigma_{j_2}(Y_n)}{2\|Y_n\|} \left(\Delta W_n^{j_1}\Delta W_n^{j_2} - \delta_{j_1,j_2}h\right)\Big\rangle. \tag{2.7}$$

The following lemmas are needed in order to prove uniform boundedness of p-th moments.

Lemma 2.2 *Let Y_n, D_n and Ω_n be given by (1.10),(2.6) and (2.5), respectively. Then*

$$1_{\Omega_n}\|Y_n\| \leq D_n, \quad \text{for all } n = 0, 1, ..., N. \tag{2.8}$$

Proof. First of all, note that $\|\Delta W_n\| \leq 1$ on Ω_{n+1} for all $0 \leq n \leq N-1$ and $N \in \mathbb{N}$. The globally Lipschitz continuity of σ and $L^{j_1}\sigma_{j_2}, j_1, j_2 \in \{1, 2, ..., m\}$, and the polynomial growth bound on μ' imply that, on $\Omega_{n+1} \cap \{\omega \in \Omega | \|Y_n(\omega)\| \leq 1\}$ for all $0 \leq n \leq N-1$,

$$\begin{aligned}
&\|Y_{n+1}\| \\
\leq& \|Y_n\| + h\|\mu(Y_n)\| + \|\sigma(Y_n)\|\|\Delta W_n\| + \frac{1}{2}\sum_{j_1,j_2=1}^{m} \|L^{j_1}\sigma_{j_2}(Y_n)\||\Delta W_n^{j_1}\Delta W_n^{j_2} - \delta_{j_1,j_2}h| \\
\leq& 1 + h\|\mu(Y_n) - \mu(0)\| + h\|\mu(0)\| + \|\sigma(Y_n) - \sigma(0)\|\|\Delta W_n\| + \|\sigma(0)\|\|\Delta W_n\| \\
&+ \frac{1}{2}\sum_{j_1,j_2=1}^{m} \left(\|L^{j_1}\sigma_{j_2}(Y_n) - L^{j_1}\sigma_{j_2}(0)\| + \|L^{j_1}\sigma_{j_2}(0)\|\right)|\Delta W_n^{j_1}\Delta W_n^{j_2} - \delta_{j_1,j_2}h| \\
\leq& 1 + hK(1 + \|Y_n\|^c)\|Y_n\| + h\|\mu(0)\| + K\|Y_n\|\|\Delta W_n\| + \|\sigma(0)\|\|\Delta W_n\| \\
&+ \frac{1}{2}\sum_{j_1,j_2=1}^{m} \left(K\|Y_n\| + \|L^{j_1}\sigma_{j_2}(0)\|\right)|\Delta W_n^{j_1}\Delta W_n^{j_2} - \delta_{j_1,j_2}h| \\
\leq& 1 + 2TK + T\|\mu(0)\| + K + \|\sigma(0)\| + \frac{m}{2}(m+T)\Big(K + \max_{1\leq j_1,j_2\leq m}\|L^{j_1}\sigma_{j_2}(0)\|\Big) \\
\leq& \lambda.
\end{aligned} \tag{2.9}$$

Moreover, the Cauchy-Schwarz inequality and the inequality $a \cdot b \leq \frac{a^2}{2} + \frac{b^2}{2}$ for all $a, b \in \mathbb{R}$ give that

$$\begin{aligned}
\|Y_{n+1}\|^2 =& \|Y_n + h\tilde{\mu}(Y_n) + \sigma(Y_n)\Delta W_n + M_n\|^2 \\
=& \|Y_n\|^2 + h^2\|\tilde{\mu}(Y_n)\|^2 + \|\sigma(Y_n)\Delta W_n\|^2 + \|M_n\|^2 \\
&+ 2h\langle Y_n, \tilde{\mu}(Y_n)\rangle + 2\langle Y_n, \sigma(Y_n)\Delta W_n\rangle + 2\langle Y_n, M_n\rangle \\
&+ 2h\langle\tilde{\mu}(Y_n), \sigma(Y_n)\Delta W_n\rangle + 2h\langle\tilde{\mu}(Y_n), M_n\rangle + 2\langle\sigma(Y_n)\Delta W_n, M_n\rangle \\
\leq& \|Y_n\|^2 + 3h^2\|\mu(Y_n)\|^2 + 3\|\sigma(Y_n)\|^2\|\Delta W_n\|^2 + 3\|M_n\|^2 \\
&+ \frac{2h}{1 + h\|\mu(Y_n)\|}\langle Y_n, \mu(Y_n)\rangle + 2\langle Y_n, \sigma(Y_n)\Delta W_n\rangle + 2\langle Y_n, M_n\rangle
\end{aligned} \tag{2.10}$$

on Ω and $0 \leq n \leq N-1$. Here we denote

$$M_n = \frac{1}{2}\sum_{j_1,j_2=1}^{m} L^{j_1}\sigma_{j_2}(Y_n)(\Delta W_n^{j_1}\Delta W_n^{j_2} - \delta_{j_1,j_2}h). \tag{2.11}$$

Additionally, the global Lipschitz continuity of $\sigma, L^{j_1}\sigma_{j_2}$ implies that for $\|x\| \geq 1$

$$\|\sigma(x)\|^2 \leq (\|\sigma(x) - \sigma(0)\| + \|\sigma(0)\|)^2 \leq (K\|x\| + \|\sigma(0)\|)^2 \leq (K + \|\sigma(0)\|)^2\|x\|^2, \tag{2.12}$$

and

$$\begin{aligned}
\|L^{j_1}\sigma_{j_2}(x)\|^2 &\leq (\|L^{j_1}\sigma_{j_2}(x) - L^{j_1}\sigma_{j_2}(0)\| + \|L^{j_1}\sigma_{j_2}(0)\|)^2 \\
&\leq (K\|x\| + \|L^{j_1}\sigma_{j_2}(0)\|)^2 \\
&\leq (K + \|L^{j_1}\sigma_{j_2}(0)\|)^2\|x\|^2,
\end{aligned} \tag{2.13}$$

and the globally one-sided Lipschitz continuity of μ gives that

$$\langle x, \mu(x)\rangle = \langle x, \mu(x) - \mu(0)\rangle + \langle x, \mu(0)\rangle \le K\|x\|^2 + \|x\| \cdot \|\mu(0)\| \le (K + \|\mu(0)\|)\|x\|^2. \tag{2.14}$$

Furthermore, the polynomial growth bound on μ' implies that

$$\begin{aligned} \|\mu(x)\|^2 &\le (\|\mu(x) - \mu(0)\| + \|\mu(0)\|)^2 \le (K(1 + \|x\|^c)\|x\| + \|\mu(0)\|)^2 \\ &\le \left(2K\|x\|^{(c+1)} + \|\mu(0)\|\right)^2 \\ &\le (2K + \|\mu(0)\|)^2 \|x\|^{2(c+1)} \\ &\le N (2K + \|\mu(0)\|)^2 \|x\|^2 \end{aligned} \tag{2.15}$$

on $1 \le \|x\| \le N^{\frac{1}{2c}}$. Combining (2.12)-(2.15), we get from (2.10) that

$$\begin{aligned} \|Y_{n+1}\|^2 \le &\|Y_n\|^2 + 3hT (2K + \|\mu(0)\|)^2 \|Y_n\|^2 + 3(K + \|\sigma(0)\|)^2\|Y_n\|^2\|\Delta W_n\|^2 \\ &+ 3\|M_n\|^2 + 2h(K + \|\mu(0)\|)\|Y_n\|^2 + 2\langle Y_n, \sigma(Y_n)\Delta W_n\rangle + 2\langle Y_n, M_n\rangle \end{aligned} \tag{2.16}$$

on $\{\omega \in \Omega | 1 \le \|Y_n(\omega)\| \le N^{\frac{1}{2c}}\}$. Since $\omega \in \Omega_{n+1}$ implies $\|\Delta W_n\| \le 1$, on $\omega \in \Omega_{n+1} \cap \{\omega \in |1 \le \|Y_n(\omega)\| \le N^{\frac{1}{2c}}\}$, we derive from (2.11) and (2.13) that

$$\begin{aligned} \|M_n\|^2 \le &\frac{m^2}{4} \sum_{j_1, j_2 = 1}^{m} \|L^{j_1}\sigma_{j_2}(Y_n)\|^2 |\Delta W_n^{j_1} \Delta W_n^{j_2} - \delta_{j_1, j_2} h|^2 \\ \le &\frac{m^2}{2} \sum_{j_1, j_2 = 1}^{m} \left(K + \|L^{j_1}\sigma_{j_2}(0)\|\right)^2 \|Y_n\|^2 \cdot \left(|\Delta W_n^{j_1} \Delta W_n^{j_2}|^2 + \delta_{j_1, j_2} h^2\right) \\ \le &\frac{m^3}{2} \Big(K + \max_{1 \le j_1, j_2 \le m} \|L^{j_1}\sigma_{j_2}(0)\|\Big)^2 \left[\|\Delta W_n\|^2 + h^2\right] \|Y_n\|^2, \end{aligned} \tag{2.17}$$

where the fact that $|\Delta W_n^{j_2}|^2 \le \|\Delta W_n\|^2 \le 1$ was used. Inserting (2.17) into (2.16) shows that

$$\begin{aligned} &\|Y_{n+1}\|^2 \\ \le &\|Y_n\|^2 \Big[1 + 3\frac{T^2}{N}(2K + \|\mu(0)\|)^2 + 3(K + \|\sigma(0)\|)^2\|\Delta W_n\|^2 + 2\frac{T}{N}(K + \|\mu(0)\|)\Big] \\ &+ \frac{3m^3}{2}\Big(K + \max_{1 \le j_1, j_2 \le m} \|L^{j_1}\sigma_{j_2}(0)\|\Big)^2 \Big(\|\Delta W_n\|^2 + h^2\Big)\|Y_n\|^2 + 2\langle Y_n, \sigma(Y_n)\Delta W_n\rangle + 2\langle Y_n, M_n\rangle \\ \le &\|Y_n\|^2 \left[1 + \frac{2}{N}\left(\frac{3}{2}(2TK + T\|\mu(0)\|)^2 + T(K + \|\mu(0)\|) + \frac{3m^3T^2}{4}\Big(K + \max_{1 \le j_1, j_2 \le m} \|L^{j_1}\sigma_{j_2}(0)\|\Big)^2\right)\right. \\ &\left. + 2\left(\frac{3}{2}(K + \|\sigma(0)\|)^2 + \frac{3m^3}{4}\Big(K + \max_{1 \le j_1, j_2 \le m} \|L^{j_1}\sigma_{j_2}(0)\|\Big)^2\right) \|\Delta W_n\|^2 + 2\alpha_n\right] \\ \le &\|Y_n\|^2 \exp\left[\frac{2\lambda}{N} + 2\lambda\|\Delta W_n\|^2 + 2\alpha_n\right] \end{aligned} \tag{2.18}$$

on $\Omega_{n+1} \cap \{\omega \in |1 \le \|Y_n(\omega)\| \le N^{\frac{1}{2c}}\}$. Now combining (2.9) and (2.18), and using mathematical induction method as used in the proof of Lemma 2.1 in [9] finish the proof. □

Lemma 2.3 *For all $p \geq 1$*

$$\sup_{\substack{N\in\mathbb{N}\\ N\geq 4\lambda pT}} \mathbb{E}\Big[\exp\Big(p\lambda\sum_{k=0}^{N-1}\|\Delta W_k\|^2\Big)\Big] < \infty. \tag{2.19}$$

Proof. This result is identical to Lemma 2.3 in [9] with only different λ. □

Lemma 2.4 *Let α_n be given by (2.7). Then for all $p \geq 1$*

$$\sup_{z\in\{-1,1\}} \sup_{N\in\mathbb{N}} \mathbb{E}\Big[\sup_{0\leq n\leq N} \exp\Big(pz\sum_{k=0}^{n-1}\alpha_k\Big)\Big] < \infty. \tag{2.20}$$

Proof. We set

$$\alpha_k^1 = 1_{\{\|Y_k\|\geq 1\}}\Big\langle \frac{Y_k}{\|Y_k\|}, \frac{\sigma(Y_k)}{\|Y_k\|}\Delta W_k\Big\rangle,$$

$$\alpha_k^2 = 1_{\{\|Y_k\|\geq 1\}}\Big\langle \frac{Y_k}{\|Y_k\|}, \sum_{j_1,j_2=1}^{m} \frac{L^{j_1}\sigma_{j_2}(Y_k)}{2\|Y_k\|}\left(\Delta W_k^{j_1}\Delta W_k^{j_2} - \delta_{j_1,j_2}h\right)\Big\rangle.$$

Then we have $\alpha_k = \alpha_k^1 + \alpha_k^2$. Hence Hölder's inequality shows that

$$\Big\|\sup_{0\leq n\leq N}\exp\Big(z\sum_{k=0}^{n-1}\alpha_k\Big)\Big\|_{L^p(\Omega;\mathbb{R})} \leq \Big\|\sup_{0\leq n\leq N}\exp\Big(z\sum_{k=0}^{n-1}\alpha_k^1\Big)\Big\|_{L^{2p}(\Omega;\mathbb{R})}\cdot\Big\|\sup_{0\leq n\leq N}\exp\Big(z\sum_{k=0}^{n-1}\alpha_k^2\Big)\Big\|_{L^{2p}(\Omega;\mathbb{R})}. \tag{2.21}$$

Note that Lemma 2.4 in [9] has proved that

$$\sup_{z\in\{-1,1\}}\Big\|\sup_{0\leq n\leq N}\exp\Big(z\sum_{k=0}^{n-1}\alpha_k^1\Big)\Big\|_{L^{2p}(\Omega;\mathbb{R})} < \infty. \tag{2.22}$$

Consequently it remains to prove the boundedness of the second term on the right-hand side of (2.21). One can easily verify that the discrete stochastic process $z\sum_{k=0}^{n-1}\alpha_k^2, n \in \{0,1,...,N\}$ is an $\{\mathcal{F}_{t_n} : 0 \leq n \leq N\}$-martingale for every $z \in \{-1,1\}$. Since the exponential function is convex, the discrete stochastic process $\exp\left(z\sum_{k=0}^{n-1}\alpha_k^2\right), n \in \{0,1,...,N\}$ is a positive $\{\mathcal{F}_{t_n} : 0 \leq n \leq N\}$-submartingale for every $z \in \{-1,1\}$. Therefore, the Doob's maximal inequality gives that

$$\Big\|\sup_{0\leq n\leq N}\exp\Big(z\sum_{k=0}^{n-1}\alpha_k^2\Big)\Big\|_{L^{2p}(\Omega;\mathbb{R})} \leq \frac{2p}{2p-1}\Big\|\exp\Big(z\sum_{k=0}^{N-1}\alpha_k^2\Big)\Big\|_{L^{2p}(\Omega;\mathbb{R})}. \tag{2.23}$$

The Cauchy-Schwarz inequality, (2.13) and Lemma 2.3 give that

$$\begin{aligned}&\Big\|\exp\Big(z\sum_{k=0}^{N-1}\alpha_k^2\Big)\Big\|_{L^{2p}(\Omega;\mathbb{R})}\\ \leq\;& \Big\|\exp\Big(\sum_{k=0}^{N-1}1_{\{\|Y_k\|\geq 1\}}\sum_{j_1,j_2=1}^{m}\frac{\|L^{j_1}\sigma_{j_2}(Y_k)\|}{2\|Y_k\|}\Big|\Delta W_k^{j_1}\Delta W_k^{j_2} - \delta_{j_1,j_2}h\Big|\Big)\Big\|_{L^{2p}(\Omega;\mathbb{R})}\end{aligned}$$

$$\begin{aligned}
&\leq \Big\| \exp\Big(\sum_{k=0}^{N-1} \big(\frac{K}{2} + \frac{1}{2} \max_{1\leq j_1, j_2 \leq m} \|L^{j_1}\sigma_{j_2}(0)\| \big) \sum_{j_1,j_2=1}^{m} \Big(\Big|\Delta W_k^{j_1} \Delta W_k^{j_2}\Big| + \Big|\delta_{j_1,j_2} h\Big| \Big) \Big) \Big\|_{L^{2p}(\Omega;\mathbb{R})} \\
&\leq \Big\| \exp\Big(\sum_{k=0}^{N-1} \big(\frac{K}{2} + \frac{1}{2} \max_{1\leq j_1, j_2 \leq m} \|L^{j_1}\sigma_{j_2}(0)\| \big) \Big(m\|\Delta W_k\|^2 + mh \Big) \Big) \Big\|_{L^{2p}(\Omega;\mathbb{R})} \\
&\leq e^{\lambda} \Big\| \exp\Big(\lambda \sum_{k=0}^{N-1} \|\Delta W_k\|^2 \Big) \Big\|_{L^{2p}(\Omega;\mathbb{R})} < \infty.
\end{aligned}$$

This together with (2.23) completes the proof. □

Lemma 2.5 *Let* D_n *be given by (2.6). Then for all* $N \geq 8\lambda p T$ *and* $p \geq 1$

$$\sup_{N\in\mathbb{N}} \mathbb{E}\left[\sup_{0\leq n\leq N} |D_n|^p \right] < \infty. \tag{2.24}$$

Proof. Note that D_n here takes the same form as D_n^N in [9], with only different α_n. With Lemma 2.3 and Lemma 2.4 at hand, one can follow the proof of Lemma 2.5 in [9] to derive the desired result. □

Lemma 2.6 *Let* Ω_N *be given by (2.5) with* $n = N$*. Then for all* $p \geq 1$

$$\sup_{N\in\mathbb{N}} \left(N^p \mathbb{P}\left[(\Omega_N)^c \right] \right) < \infty. \tag{2.25}$$

Proof. The proof is identical to the proof of Lemma 2.6 in [9]. □

Before establishing the main result of this section, we also need two Burkholder-Davis-Gundy type inequalities.

Lemma 2.7 *Let* $k \in \mathbb{N}$ *and let* $Z : [0,T] \times \Omega \to \mathbb{R}^{k\times m}$ *be a predictable stochastic process satisfying* $\mathbb{P}(\int_0^T \|Z_s\|^2 ds < \infty) = 1$*. Then for all* $t \in [0,T]$ *and all* $p \geq 2$

$$\Big\| \sup_{s\in[0,t]} \Big\| \int_0^s Z_u dW_u \Big\| \Big\|_{L^p(\Omega;\mathbb{R})} \leq p \Big(\int_0^t \sum_{i=1}^{m} \|Z_s e_i\|^2_{L^p(\Omega;\mathbb{R}^k)} ds \Big)^{\frac{1}{2}}. \tag{2.26}$$

Here the vectors $e_1 = (1,0,...,0)^T \in \mathbb{R}^m$*,* $e_2 = (0,1,...,0)^T \in \mathbb{R}^m$*, ...,* $e_m = (0,0,...,1)^T \in \mathbb{R}^m$ *are orthogonal basis of vector space* $\mathbb{R}^m$*.*

Proof. Combining Doob's maximal inequality and Lemma 7.7 of Da Prato, G., and Zabczyk [2] gives the desired assertion. □

The following is a discrete version of the Burkholder-Davis-Gundy type inequality (2.26).

Lemma 2.8 *Let* $k \in \mathbb{N}$ *and let* $Z_l : \Omega \to \mathbb{R}^{k\times m}, l \in \{0,1,...,N-1\}$ *be a family of mappings such that* Z_l *is* $\mathcal{F}_{\frac{lT}{N}}/\mathcal{B}(\mathbb{R}^{k\times m})$*-measurable. Then for all* $0 \leq n \leq N$ *and* $p \geq 2$

$$\Big\| \sup_{0\leq j\leq n} \Big\| \sum_{l=0}^{j-1} Z_l \Delta W_l \Big\| \Big\|_{L^p(\Omega;\mathbb{R})} \leq p \Big(\sum_{l=0}^{n-1} \sum_{i=1}^{m} \|Z_l e_i\|^2_{L^p(\Omega;\mathbb{R}^k)} \frac{T}{N} \Big)^{\frac{1}{2}}. \tag{2.27}$$

Theorem 2.9 *Let Y_n be given by (1.10). Then for all $p \in [1, \infty)$*

$$\sup_{N \in \mathbb{N}} \left[\sup_{0 \le n \le N} \mathbb{E} \|Y_n\|^p \right] < \infty. \tag{2.28}$$

Proof. From (1.10) we have

$$\begin{aligned} \|Y_n\|_{L^p(\Omega;\mathbb{R}^d)} \le & \|\xi\|_{L^p(\Omega;\mathbb{R}^d)} \\ & + \Big\| \sum_{k=0}^{n-1} h\tilde{\mu}(Y_k) \Big\|_{L^p(\Omega;\mathbb{R}^d)} + \Big\| \sum_{k=0}^{n-1} \sigma(Y_k)\Delta W_k \Big\|_{L^p(\Omega;\mathbb{R}^d)} + \Big\| \sum_{k=0}^{n-1} M_k \Big\|_{L^p(\Omega;\mathbb{R}^d)}, \end{aligned} \tag{2.29}$$

where the notation M_k comes from (2.11). Using (2.2), the triangle inequality and the Burkholder-Davis-Gundy type inequality in Lemma 2.8 we have

$$\begin{aligned} \Big\| \sum_{k=0}^{n-1} \sigma(Y_k)\Delta W_k \Big\|_{L^p(\Omega;\mathbb{R}^d)} \le & \Big\| \sum_{k=0}^{n-1} \left[\sigma(Y_k) - \sigma(0)\right] \Delta W_k \Big\|_{L^p(\Omega;\mathbb{R}^d)} + \Big\| \sum_{k=0}^{n-1} \sigma(0)\Delta W_k \Big\|_{L^p(\Omega;\mathbb{R}^d)} \\ \le & p\Big(\sum_{k=0}^{n-1} \sum_{i=1}^{m} \|\sigma_i(Y_k) - \sigma_i(0)\|^2_{L^p(\Omega;\mathbb{R}^d)} h \Big)^{1/2} + p\Big(\frac{nT}{N} \sum_{i=1}^{m} \|\sigma_i(0)\|^2 \Big)^{1/2} \\ \le & p\Big(mK^2 h \sum_{k=0}^{n-1} \|Y_k\|^2_{L^p(\Omega;\mathbb{R}^d)} \Big)^{1/2} + p\sqrt{Tm}\|\sigma(0)\|. \end{aligned} \tag{2.30}$$

For the fourth term on the right-hand side of (2.29), the estimate in the second inequality of (2.13) and the independence of Y_k and ΔW_k imply that

$$\begin{aligned} \Big\| \sum_{k=0}^{n-1} M_k \Big\|_{L^p(\Omega;\mathbb{R}^d)} = & \frac{1}{2} \Big\| \sum_{k=0}^{n-1} \Big(\sum_{j_1,j_2=1}^{m} L^{j_1}\sigma_{j_2}(Y_k) \left(\Delta W_k^{j_1} \Delta W_k^{j_2} - \delta_{j_1,j_2} h \right) \Big) \Big\|_{L^p(\Omega;\mathbb{R}^d)} \\ \le & \frac{1}{2} \sum_{k=0}^{n-1} \sum_{j_1,j_2=1}^{m} \Big[\Big(K \|Y_k\|_{L^p(\Omega;\mathbb{R}^d)} + \|L^{j_1}\sigma_{j_2}(0)\| \Big) \cdot \Big\| \Delta W_k^{j_1} \Delta W_k^{j_2} - \delta_{j_1,j_2} h \Big\|_{L^p(\Omega;\mathbb{R})} \Big]. \end{aligned} \tag{2.31}$$

Using Burkholder-Davis-Gundy type inequality (2.27) and mutual independence of ΔW_k gives

$$\begin{aligned} & \frac{1}{2} \sum_{j_1,j_2=1}^{m} \|\Delta W_k^{j_1} \Delta W_k^{j_2} - \delta_{j_1,j_2} h\|_{L^p(\Omega;\mathbb{R})} \le \frac{1}{2} \sum_{j_1,j_2=1}^{m} \|\Delta W_k^{j_1} \Delta W_k^{j_2}\|_{L^p(\Omega;\mathbb{R})} + \frac{1}{2} mh \\ = & \frac{1}{2} \sum_{\substack{j_1,j_2=1 \\ j_1 \ne j_2}}^{m} \|\Delta W_k^{j_1}\|_{L^p(\Omega;\mathbb{R})} \cdot \|\Delta W_k^{j_2}\|_{L^p(\Omega;\mathbb{R})} + \frac{1}{2} \sum_{j=1}^{m} \|\Delta W_k^{j}\|^2_{L^{2p}(\Omega;\mathbb{R})} + \frac{1}{2} mh \\ \le & c_{p,m} h, \end{aligned} \tag{2.32}$$

where $c_{p,m} = \frac{m^2 - m}{2} p^2 + 2mp^2 + \frac{m}{2}$. Inserting (2.32) into (2.31) we obtain that

$$\Big\| \sum_{k=0}^{n-1} M_k \Big\|_{L^p(\Omega;\mathbb{R}^d)} \le K c_{p,m} h \sum_{k=0}^{n-1} \|Y_k\|_{L^p(\Omega;\mathbb{R}^d)} + T c_{p,m} \sup_{1 \le j_1, j_2 \le m} \|L^{j_1}\sigma_{j_2}(0)\|. \tag{2.33}$$

Combining (2.29),(2.30) and (2.33), and using $\|\tilde{\mu}(Y_k)\| \leq 1$ give

$$\begin{aligned}\|Y_n\|_{L^p(\Omega;\mathbb{R}^d)} \leq & \|\xi\|_{L^p(\Omega;\mathbb{R}^d)} + N + p\Big(mK^2h\sum_{k=0}^{n-1}\|Y_k\|^2_{L^p(\Omega;\mathbb{R}^d)}\Big)^{1/2} + p\sqrt{Tm}\|\sigma(0)\| \\ & + Kc_{p,m}h\sum_{k=0}^{n-1}\|Y_k\|_{L^p(\Omega;\mathbb{R}^d)} + Tc_{p,m}\sup_{1\leq j_1,j_2\leq m}\|L^{j_1}\sigma_{j_2}(0)\|.\end{aligned} \tag{2.34}$$

Thus taking square of both sides shows that

$$\begin{aligned}\|Y_n\|^2_{L^p(\Omega;\mathbb{R}^d)} \leq & 3\Big(\|\xi\|_{L^p(\Omega;\mathbb{R}^d)} + N + p\sqrt{Tm}\|\sigma(0)\| + Tc_{p,m}\sup_{1\leq j_1,j_2\leq m}\|L^{j_1}\sigma_{j_2}(0)\|\Big)^2 \\ & + 3mK^2p^2h\sum_{k=0}^{n-1}\|Y_k\|^2_{L^p(\Omega;\mathbb{R}^d)} + 3K^2c^2_{m,p}Th\sum_{k=0}^{n-1}\|Y_k\|^2_{L^p(\Omega;\mathbb{R}^d)}.\end{aligned} \tag{2.35}$$

In the next step Gronwall's lemma gives that

$$\sup_{0\leq n\leq N}\|Y_n\|_{L^p(\Omega;\mathbb{R}^d)} \leq \sqrt{3}\exp(C_1)\left(\|\xi\|_{L^p(\Omega;\mathbb{R}^d)} + N + C_2\right), \tag{2.36}$$

where $C_1 = \frac{1}{2}(3mK^2p^2 + 3K^2c^2_{p,m}T)T$, $C_2 = p\sqrt{Tm}\|\sigma(0)\| + Tc_{p,m}\sup_{1\leq j_1,j_2\leq m}\|L^{j_1}\sigma_{j_2}(0)\|$. Due to the N on the right-hand side of (2.36), (2.36) does not complete the prove. However, exploiting (2.36) in an appropriate bootstrap argument will enable us to establish (2.28). First, Hölder's inequality, Lemma 2.6 and the estimate (2.36) show that

$$\begin{aligned}& \sup_{N\in\mathbb{N}}\sup_{0\leq n\leq N}\|1_{(\Omega_n)^c}Y_n\|_{L^p(\Omega;\mathbb{R}^d)} \\ \leq & \sup_{N\in\mathbb{N}}\sup_{0\leq n\leq N}\left(\|1_{(\Omega_n)^c}\|_{L^{2p}(\Omega;\mathbb{R}^d)}\|Y_n\|_{L^{2p}(\Omega;\mathbb{R}^d)}\right) \\ \leq & \Big(\sup_{N\in\mathbb{N}}\left(N\cdot\|1_{(\Omega_N)^c}\|_{L^{2p}(\Omega;\mathbb{R}^d)}\right)\Big)\Big(\sup_{N\in\mathbb{N}}\sup_{0\leq n\leq N}\left(N^{-1}\cdot\|Y_n\|_{L^{2p}(\Omega;\mathbb{R}^d)}\right)\Big) \\ \leq & \Big(\sup_{N\in\mathbb{N}}N^{2p}\cdot\mathbb{P}\left[(\Omega_N)^c\right]\Big)^{\frac{1}{2p}}\Big(\sup_{N\in\mathbb{N}}\sup_{0\leq n\leq N}\left(N^{-1}\cdot\|Y_n\|_{L^{2p}(\Omega;\mathbb{R}^d)}\right)\Big) \\ < & \infty.\end{aligned} \tag{2.37}$$

In addition, Lemma 2.2 and Lemma 2.5 imply that

$$\sup_{N\in\mathbb{N}}\sup_{0\leq n\leq N}\|1_{\Omega_n}Y_n\|_{L^p(\Omega;\mathbb{R}^d)} \leq \sup_{N\in\mathbb{N}}\sup_{0\leq n\leq N}\|D_n\|_{L^p(\Omega;\mathbb{R}^d)} < \infty. \tag{2.38}$$

Combining (2.37) and (2.38) finally completes the proof. □

3 Strong convergence order of the tamed Milstein method

In order to recover the strong convergence order one for the tamed Milstein method, we additionally need the following assumptions. Throughout this section $C_{p,T}$ is a generic constant that might vary from one place to another and depends on μ, σ, the initial data ξ, and the interval of integration $[0, T]$, but is independent of the discretisation parameter.

Assumption 3.1 *Assume that $\mu(x)$ and $\sigma_i(x)$ are two times continuous differentiable and there exist positive constants $K, q \geq 1$, such that $\forall x \in \mathbb{R}^d, i = 1, ..., d$*

$$\begin{aligned} \|\mu''(x)\|_{L^{(2)}(\mathbb{R}^d;\mathbb{R}^d)} &\leq K(1 + \|x\|^q), && (3.1)\\ \|\sigma_i''(x)\|_{L^{(2)}(\mathbb{R}^d;\mathbb{R}^d)} &\leq K. && (3.2) \end{aligned}$$

Here, for a two times continuous differentiable function $f : \mathbb{R}^d \to \mathbb{R}^d$ we use the notation $\|f''(x)\|_{L^{(2)}(\mathbb{R}^d;\mathbb{R}^d)} = \sup_{h_1,h_2 \in \mathbb{R}^d, \|h_1\| \leq 1, \|h_2\| \leq 1} \|f''(x)(h_1, h_2)\|$. The bilinear operator $f''(x) : \mathbb{R}^d \times \mathbb{R}^d \to \mathbb{R}^d$ is defined by (3.8). In the following convergence analysis, we prefer to write the tamed Milstein method in the form of (1.8), rather than (1.10). We now introduce appropriate time continuous interpolations of the time discrete numerical approximations. More accurately, we define the time continuous approximation $\bar{Y}_s$ such that for $s \in [t_n, t_{n+1})$

$$\begin{aligned} \bar{Y}_s :=& Y_n + (s - t_n)\tilde{\mu}(Y_n) + \sigma(Y_n)(W_s - W_{t_n}) + \sum_{j_1,j_2=1}^{m} L^{j_1}\sigma_{j_2}(Y_n) I_{j_1,j_2}^{t_n,s} \\ =& Y_n + \int_{t_n}^{s} \tilde{\mu}(Y_n)du + \sum_{i=1}^{m} \int_{t_n}^{s} \sigma_i(Y_n)dW_u^i + \sum_{i=1}^{m} \int_{t_n}^{s} \sum_{j=1}^{m} L^j\sigma_i(Y_n)\Delta W_u^j dW_u^i, \end{aligned} \tag{3.3}$$

where we use the notation

$$I_{j_1,j_2}^{t_n,s} = \int_{t_n}^{s} \int_{t_n}^{s_2} dW_{s_1}^{j_1} dW_{s_2}^{j_2}, \quad \Delta W_s^j := \sum_{n=0}^{\infty} 1_{\{t_n \leq s < t_{n+1}\}}(W_s^j - W_{t_n}^j).$$

It is evident that $\bar{Y}_{t_n} = Y_n, n = 0, 1, ..., N$, that is, $\bar{Y}_t$ coincides with the discrete solutions at the grid-points. We can rewrite $\bar{Y}_t$ as an integral form in the whole interval $[0, T]$

$$\bar{Y}_t = Y_0 + \int_0^t \tilde{\mu}(Y_{n_s})ds + \sum_{i=1}^{m} \int_0^t \Big[\sigma_i(Y_{n_s}) + \sum_{j=1}^{m} L^j\sigma_i(Y_{n_s})\Delta W_s^j\Big] dW_s^i, \tag{3.4}$$

where n_s is the greatest integer number such that $t_{n_s} \leq s$. Combining (1.2) and (3.4) gives

$$X_t - \bar{Y}_t = \int_0^t [\mu(X_s) - \tilde{\mu}(Y_{n_s})] ds + \sum_{i=1}^{m} \int_0^t \Big[\sigma_i(X_s) - \sigma_i(Y_{n_s}) - \sum_{j=1}^{m} L^j\sigma_i(Y_{n_s})\Delta W_s^j\Big] dW_s^i. \tag{3.5}$$

In what follows, we also use deterministic Taylor formula frequently. If a function $f : \mathbb{R}^d \to \mathbb{R}^d$ is twice differentiable, the following Taylor's formula holds [1]

$$f(\bar{Y}_s) - f(Y_{n_s}) = f'(Y_{n_s})(\bar{Y}_s - Y_{n_s}) + R_1(f), \tag{3.6}$$

where $R_1(f)$ is the remainder term

$$R_1(f) = \int_0^1 (1 - r) f''(Y_{n_s} + r(\bar{Y}_s - Y_{n_s}))(\bar{Y}_s - Y_{n_s}, \bar{Y}_s - Y_{n_s})dr. \tag{3.7}$$

Here for arbitrary $a, h_1, h_2 \in \mathbb{R}^d$ the derivatives have the following expression

$$f'(a)(h_1) = \sum_{i=1}^{d} \frac{\partial f}{\partial x^i} h_1^i, \quad f''(a)(h_1, h_2) = \sum_{i,j=1}^{d} \frac{\partial^2 f}{\partial x^i \partial x^j} h_1^i h_2^j. \tag{3.8}$$

Replacing $\bar{Y}_s - Y_{n_s}$ in (3.6) with (3.3) and rearranging lead to

$$f(\bar{Y}_s) - f(Y_{n_s}) = f'(Y_{n_s})\big(\sigma(Y_{n_s})(W_s - W_{t_{n_s}})\big) + \tilde{R}_1(f), \tag{3.9}$$

where

$$\tilde{R}_1(f) = f'(Y_{n_s})\Big((s - t_{n_s})\tilde{\mu}(Y_{n_s}) + \sum_{j_1,j_2=1}^{m} L^{j_1}\sigma_{j_2}(Y_{n_s}) I_{j_1,j_2}^{t_{n_s},s}\Big) + R_1(f). \tag{3.10}$$

By the definitions (1.7) and (3.8), it can be readily checked that

$$\sigma_i'(x)\big(\sigma_j(x)\big) = L^j \sigma_i(x). \tag{3.11}$$

Therefore replacing f in (3.9) by σ_i and taking (3.11) into account show that

$$\tilde{R}_1(\sigma_i) = \sigma_i(\bar{Y}_s) - \sigma_i(Y_{n_s}) - \sum_{j=1}^{m} L^j \sigma_i(Y_{n_s}) \Delta W_s^j. \tag{3.12}$$

Theorem 3.2 *Let conditions in Assumptions 2.1 and 3.1 and (1.9) be fulfilled. Then there exists a family of real numbers* $C_{p,T} \geq 1$, $p \geq 1$ *such that*

$$\left(\mathbb{E}\Big[\sup_{t \in [0,T]} \|X_t - \bar{Y}_t\|^p\Big]\right)^{1/p} \leq C_{p,T} \cdot h, \quad h \in (0, 1]. \tag{3.13}$$

The following five lemmas are used in the proof of Theorem 3.2.

Lemma 3.3 *Let conditions in Assumptions 2.1 be fulfilled. Then for all* $p \geq 1, 1 \leq j_1, j_2 \leq m$ *the following estimates hold*

$$\sup_{N \in \mathbb{N}} \sup_{0 \leq n \leq N} \Big[\mathbb{E}\|\mu(Y_n)\|^p \bigvee \mathbb{E}\|\mu'(Y_n)\|^p \bigvee \mathbb{E}\|\sigma(Y_n)\|^p \bigvee \mathbb{E}\|L^{j_1}\sigma_{j_2}(Y_n)\|^p\Big] < \infty. \tag{3.14}$$

Proof. It immediately follows from Theorem 2.9 by considering (2.4) and (2.12)-(2.15). □

Lemma 3.4 *Let conditions in Theorem 3.2 be fulfilled. Then for all* $p \geq 1$

$$\sup_{t \in [0,T]} \Big[\|X_t\|_{L^p(\Omega;\mathbb{R}^d)} \bigvee \|\mu(X_t)\|_{L^p(\Omega;\mathbb{R}^d)} \bigvee \|\sigma(X_t)\|_{L^p(\Omega;\mathbb{R}^{d\times m})} \bigvee \|\bar{Y}_t\|_{L^p(\Omega;\mathbb{R}^d)}\Big] < \infty. \tag{3.15}$$

Proof. Conditions (2.1)-(2.2) ensure that the exact solution X_t satisfies $\sup_{t\in[0,T]} \mathbb{E}\|X_t\|^p < \infty$ (See, e.g., [14, Theorem 4.1]). Polynomial growth condition on μ as (2.15) and linear growth condition on σ give the second and the third estimates. Taking the definition (3.3) and Lemma 3.3 into account, we can easily obtain the last estimate. □

Lemma 3.5 *Let conditions in Theorem 3.2 be fulfilled. Then there exists a family of constants* $C_{p,T} \geq 1$ *such that for* $p \geq 1$

$$\sup_{t\in[0,T]} \|\bar{Y}_t - Y_{n_t}\|_{L^p(\Omega;\mathbb{R}^d)} \bigvee \sup_{t\in[0,T]} \|\mu(X_t) - \mu(X_{t_{n_t}})\|_{L^p(\Omega;\mathbb{R}^d)} \leq C_{p,T}\, h^{\frac{1}{2}}. \tag{3.16}$$

Proof. Let n_t be the greatest integer number such that $t_{n_t} \leq t$. From (3.3) we have

$$\bar{Y}_t - Y_{n_t} = (t - t_{n_t})\tilde{\mu}(Y_{n_t}) + \sigma(Y_{n_t})(W_t - W_{t_{n_t}}) + \sum_{j_1,j_2=1}^{m} L^{j_1}\sigma_{j_2}(Y_{n_t}) I_{j_1,j_2}^{t_{n_t},t}. \tag{3.17}$$

Following the same line as estimating (2.29), one can readily derive the first estimate. For the second estimate, we use (2.4) and Hölder's inequality to obtain

$$\begin{aligned}\|\mu(X_t) - \mu(X_{t_{n_t}})\|_{L^p(\Omega;\mathbb{R}^d)} \leq & K\big\|(1 + \|X_t\|^c + \|X_{t_{n_t}}\|^c) \cdot \|X_t - X_{t_{n_t}}\|\big\|_{L^p(\Omega;\mathbb{R})} \\ \leq & K\big\|(1 + \|X_t\|^c + \|X_{t_{n_t}}\|^c)\big\|_{L^{2p}(\Omega;\mathbb{R})} \cdot \big\|X_t - X_{t_{n_t}}\big\|_{L^{2p}(\Omega;\mathbb{R}^d)},\end{aligned} \tag{3.18}$$

where

$$X_t - X_{t_{n_t}} = \int_{t_{n_t}}^{t} \mu(X_s)ds + \int_{t_{n_t}}^{t} \sigma(X_s)dW_s. \tag{3.19}$$

Taking Lemma 3.4 into account and using the same argument as in the previous section, one can obtain $\big\|X_t - X_{t_{n_t}}\big\|_{L^{2p}(\Omega;\mathbb{R}^d)} \leq C_{p,T}h^{\frac{1}{2}}$, and then complete the proof easily by considering (3.18) and Lemma 3.4. □

Lemma 3.6 *Let* $p \geq 1$ *and let conditions in Theorem 3.2 be fulfilled. Then for* $i = 1, 2, ..., m$

$$\|\tilde{R}_1(\mu)\|_{L^p(\Omega;\mathbb{R}^d)} \vee \|\tilde{R}_1(\sigma_i)\|_{L^p(\Omega;\mathbb{R}^d)} \leq C_{p,T}h. \tag{3.20}$$

Proof. To estimate $\|\tilde{R}_1(\mu)\|_{L^p(\Omega;\mathbb{R}^d)}$, we need to estimate $\|R_1(\mu)\|_{L^p(\Omega;\mathbb{R}^d)}$ first. Due to Theorem 2.9 and Lemma 3.4 we can find some suitable constant $C_{p,T}$ such that

$$\begin{aligned}\|R_1(\mu)\|_{L^p(\Omega;\mathbb{R}^d)} \leq & \int_0^1 (1-r)\Big\|\mu^{''}(Y_{n_s} + r(\bar{Y}_s - Y_{n_s}))\, \big(\bar{Y}_s - Y_{n_s}, \bar{Y}_s - Y_{n_s}\big)\Big\|_{L^p(\Omega;\mathbb{R})} dr \\ \leq & \int_0^1 \Big\|\|\mu^{''}(Y_{n_s} + r(\bar{Y}_s - Y_{n_s}))\|_{L^{(2)}(\mathbb{R}^d;\mathbb{R}^d)} \cdot \|\bar{Y}_s - Y_{n_s}\|^2\Big\|_{L^p(\Omega;\mathbb{R})} dr \\ \leq & K\left\|\big(1 + \|Y_{n_s}\|^q + \|\bar{Y}_s\|^q\big) \cdot \|\bar{Y}_s - Y_{n_s}\|^2\right\|_{L^p(\Omega;\mathbb{R})} \\ \leq & K\left\|1 + \|Y_{n_s}\|^q + \|\bar{Y}_s\|^q\right\|_{L^{2p}(\Omega;\mathbb{R})} \cdot \left\|\bar{Y}_s - Y_{n_s}\right\|_{L^{4p}(\Omega;\mathbb{R}^d)}^2 \\ \leq & C_{p,T}h,\end{aligned} \tag{3.21}$$

where the polynomial growth condition (3.1) on $\mu''(x)$, Lemma 3.5, Hölder's inequality and Jensen's inequality were also used. Now we return to $\|\tilde{R}_1(\mu)\|_{L^p(\Omega;\mathbb{R}^d)}$. Replacing f in (3.10) with μ gives

$$\|\tilde{R}_1(\mu)\|_{L^p(\Omega;\mathbb{R}^d)} \tag{3.22}$$

$$
\begin{aligned}
&\leq \|\mu'(Y_{n_s})(s-t_{n_s})\tilde{\mu}(Y_{n_s})\|_{L^p(\Omega;\mathbb{R}^d)} + \Big\|\mu'(Y_{n_s})\sum_{j_1,j_2=1}^{m} L^{j_1}\sigma_{j_2}(Y_{n_s})I_{j_1,j_2}^{t_{n_s},s}\Big\|_{L^p(\Omega;\mathbb{R}^d)} + \|R_1(\mu)\|_{L^p(\Omega;\mathbb{R}^d)}\\
&\leq h\,\|\mu'(Y_{n_s})\mu(Y_{n_s})\|_{L^p(\Omega;\mathbb{R}^d)} + \frac{1}{2}\sum_{j_1,j_2=1}^{m}\Big\|\mu'(Y_{n_s})\Big(L^{j_1}\sigma_{j_2}(Y_{n_s})(\Delta W_s^{j_1}\Delta W_s^{j_2}-\delta_{j_1,j_2}h)\Big)\Big\|_{L^p(\Omega;\mathbb{R}^d)}\\
&\quad + \|R_1(\mu)\|_{L^p(\Omega;\mathbb{R}^d)}.
\end{aligned}
$$

Following the same line as estimating (2.32) and noticing that $s - t_{n_s} \leq h$, one can similarly arrive at

$$
\frac{1}{2}\sum_{j_1,j_2=1}^{m}\|\Delta W_s^{j_1}\Delta W_s^{j_2}-\delta_{j_1,j_2}h\|_{L^p(\Omega;\mathbb{R})} \leq c_{p,m}h. \tag{3.23}
$$

Further, using Hölder's inequality we derive from Lemma 3.3 that for $0 \leq s \leq t \leq T, 1 \leq j_1, j_2 \leq m$

$$
\begin{aligned}
\|\mu'(Y_{n_s})\mu(Y_{n_s})\|_{L^p(\Omega;\mathbb{R}^d)} &\leq \|\mu'(Y_{n_s})\|_{L^{2p}(\Omega;\mathbb{R}^{d\times d})}\cdot\|\mu(Y_{n_s})\|_{L^{2p}(\Omega;\mathbb{R}^d)} < \infty,\\
\big\|\mu'(Y_{n_s})L^{j_1}\sigma_{j_2}(Y_{n_s})\big\|_{L^p(\Omega;\mathbb{R}^d)} &\leq \|\mu'(Y_{n_s})\|_{L^{2p}(\Omega;\mathbb{R}^{d\times d})}\cdot\big\|L^{j_1}\sigma_{j_2}(Y_{n_s})\big\|_{L^{2p}(\Omega;\mathbb{R}^d)} < \infty.
\end{aligned} \tag{3.24}
$$

Now, using the independence of Y_{n_s} and $\Delta W_s^{j_1}, \Delta W_s^{j_2}$, and combining (3.21), (3.22), (3.23) and (3.24), one can show

$$
\|\tilde{R}_1(\mu)\|_{L^p(\Omega;\mathbb{R}^d)} \leq C_{p,T}h. \tag{3.25}
$$

In a similar way as estimating $\|\tilde{R}_1(\mu)\|_{L^p(\Omega;\mathbb{R}^d)}$, one can derive that

$$
\|\tilde{R}_1(\sigma_i)\|_{L^p(\Omega;\mathbb{R}^d)} \leq C_{p,T}h. \quad \square \tag{3.26}
$$

Our proof of Theorem 3.2 also needs the following Burkholder-Davis-Gundy type inequality for discrete-time martingale (see Theorem 3.28 in [11] and Lemma 5.1 in [8]).

Lemma 3.7 *Let $Z_1, ..., Z_N : \Omega \to \mathbb{R}$ be $\mathcal{F}/\mathcal{B}(\mathbb{R})$-measurable mappings with $\mathbb{E}\|Z_n\|^p < \infty$ for all $n \in \{1, ..., N\}$ and with $\mathbb{E}[Z_{n+1}|Z_1, ..., Z_n] = 0$ for all $n \in \{1, ..., N\}$. Then*

$$
\|Z_1 + ... + Z_n\|_{L^p(\Omega;\mathbb{R})} \leq c_p\left(\|Z_1\|^2_{L^p(\Omega;\mathbb{R})} + ... + \|Z_n\|^2_{L^p(\Omega;\mathbb{R})}\right)^{\frac{1}{2}} \tag{3.27}
$$

for every $p \in [2, \infty)$, where c_p are constants dependent of p, but independent of n.

Proof of Theorem 3.2. Applying Itô's formula to (3.5) gives

$$
\begin{aligned}
\|X_s - \bar{Y}_s\|^2 =& 2\int_0^s \Big\langle X_u - \bar{Y}_u, \mu(X_u) - \tilde{\mu}(Y_{n_u})\Big\rangle du\\
&+ 2\sum_{i=1}^{m}\int_0^s \Big\langle X_u - \bar{Y}_u, \sigma_i(X_u) - \sigma_i(Y_{n_u}) - \sum_{j=1}^{m} L^j\sigma_i(Y_{n_u})\Delta W_u^j\Big\rangle dW_u^i\\
&+ \sum_{i=1}^{m}\int_0^s \Big\|\sigma_i(X_u) - \sigma_i(Y_{n_u}) - \sum_{j=1}^{m} L^j\sigma_i(Y_{n_u})\Delta W_u^j\Big\|^2 du.
\end{aligned} \tag{3.28}
$$

For the integrand of the first term in (3.28), we use (2.1) and the Cauchy-Schwarz inequality to arrive at

$$\begin{aligned}
&\Big\langle X_u - \bar{Y}_u, \mu(X_u) - \tilde{\mu}(Y_{n_u})\Big\rangle \\
= \;& \Big\langle X_u - \bar{Y}_u, \mu(X_u) - \mu(\bar{Y}_u)\Big\rangle + \Big\langle X_u - \bar{Y}_u, \mu(\bar{Y}_u) - \mu(Y_{n_u})\Big\rangle + \Big\langle X_u - \bar{Y}_u, h\, \|\mu(Y_{n_u})\|\, \tilde{\mu}(Y_{n_u})\Big\rangle \\
\leq \;& K\|X_u - \bar{Y}_u\|^2 + \Big\langle X_u - \bar{Y}_u, \mu(\bar{Y}_u) - \mu(Y_{n_u})\Big\rangle + \frac{1}{2}\|X_u - \bar{Y}_u\|^2 + \frac{1}{2}h^2\|\mu(Y_{n_u})\|^2\, \|\tilde{\mu}(Y_{n_u})\|^2 \\
\leq \;& (K + \frac{1}{2})\|X_u - \bar{Y}_u\|^2 + \Big\langle X_u - \bar{Y}_u, \mu(\bar{Y}_u) - \mu(Y_{n_u})\Big\rangle + \frac{1}{2}h^2\|\mu(Y_{n_u})\|^4. \qquad (3.29)
\end{aligned}$$

For the integrand of the third term in (3.28), one can use an elementary inequality, the notation (3.12) and (2.2) to get

$$\begin{aligned}
\Big\|\sigma_i(X_u) - \sigma_i(Y_{n_u}) - \sum_{j=1}^{m} L^j\sigma_i(Y_{n_u})\Delta W_u^j\Big\|^2 \leq & 2\|\sigma_i(X_u) - \sigma_i(\bar{Y}_u)\|^2 + 2\|\tilde{R}_1(\sigma_i)\|^2 \\
\leq & 2K^2\|X_u - \bar{Y}_u\|^2 + 2\|\tilde{R}_1(\sigma_i)\|^2.
\end{aligned} \qquad (3.30)$$

Inserting (3.29) and (3.30) into (3.28) and using the notation (3.12) show

$$\begin{aligned}
\|X_s - \bar{Y}_s\|^2 = & (2K + 1 + 2mK^2)\int_0^s \|X_u - \bar{Y}_u\|^2 du + h^2 \int_0^s \|\mu(Y_{n_u})\|^4 du \\
& + 2\sum_{i=1}^{m}\int_0^s \|\tilde{R}_1(\sigma_i)\|^2 du + 2\int_0^s \Big\langle X_u - \bar{Y}_u, \mu(\bar{Y}_u) - \mu(Y_{n_u})\Big\rangle du \\
& + 2\sum_{i=1}^{m}\int_0^s \Big\langle X_u - \bar{Y}_u, \sigma_i(X_u) - \sigma_i(\bar{Y}_u) + \tilde{R}_1(\sigma_i)\Big\rangle dW_u^i.
\end{aligned} \qquad (3.31)$$

Hence

$$\begin{aligned}
\sup_{0\leq s\leq t}\|X_s - \bar{Y}_s\|^2 \leq & (2K + 1 + 2mK^2)\int_0^t \|X_s - \bar{Y}_s\|^2 ds + h^2 \int_0^t \|\mu(Y_{n_s})\|^4 ds \\
& + 2\sum_{i=1}^{m}\int_0^t \|\tilde{R}_1(\sigma_i)\|^2 ds + 2\sup_{0\leq s\leq t}\int_0^s \Big\langle X_u - \bar{Y}_u, \mu(\bar{Y}_u) - \mu(Y_{n_u})\Big\rangle du \\
& + 2\sup_{0\leq s\leq t}\sum_{i=1}^{m}\int_0^s \Big\langle X_u - \bar{Y}_u, \sigma_i(X_u) - \sigma_i(\bar{Y}_u) + \tilde{R}_1(\sigma_i)\Big\rangle dW_u^i,
\end{aligned}$$

and thus for $p \geq 4$

$$\begin{aligned}
&\Big\|\sup_{0\leq s\leq t}\|X_s - \bar{Y}_s\|\Big\|^2_{L^p(\Omega;\mathbb{R})} = \Big\|\sup_{0\leq s\leq t}\|X_s - \bar{Y}_s\|^2\Big\|_{L^{\frac{p}{2}}(\Omega;\mathbb{R})} \\
\leq & (2K + 1 + 2mK^2)\int_0^t \|X_s - \bar{Y}_s\|^2_{L^p(\Omega;\mathbb{R}^d)} ds + h^2 \int_0^t \|\mu(Y_{n_s})\|^4_{L^{2p}(\Omega;\mathbb{R}^d)} ds \\
& + 2\sum_{i=1}^{m}\int_0^t \|\tilde{R}_1(\sigma_i)\|^2_{L^p(\Omega;\mathbb{R}^d)} ds + 2\Big\|\sup_{0\leq s\leq t}\int_0^s \Big\langle X_u - \bar{Y}_u, \mu(\bar{Y}_u) - \mu(Y_{n_u})\Big\rangle du\Big\|_{L^{\frac{p}{2}}(\Omega;\mathbb{R})} \\
& + 2\Big\|\sup_{0\leq s\leq t}\sum_{i=1}^{m}\int_0^s \Big\langle X_u - \bar{Y}_u, \sigma_i(X_u) - \sigma_i(\bar{Y}_u) + \tilde{R}_1(\sigma_i)\Big\rangle dW_u^i\Big\|_{L^{\frac{p}{2}}(\Omega;\mathbb{R})}.
\end{aligned} \qquad (3.32)$$

For the last term on the right-hand side of (3.32), we use (2.2), (2.26), the Cauchy-Schwarz inequality and elementary inequalities to derive

$$
\begin{aligned}
&2\Big\| \sup_{0\le s\le t} \sum_{i=1}^{m} \int_0^s \Big\langle X_u-\bar{Y}_u, \sigma_i(X_u)-\sigma_i(\bar{Y}_u)+\tilde{R}_1(\sigma_i)\Big\rangle dW_u^i \Big\|_{L^{\frac{p}{2}}(\Omega;\mathbb{R})}\\
\le\;& 2\sum_{i=1}^{m} \Big\| \sup_{0\le s\le t} \int_0^s \Big\langle X_u-\bar{Y}_u, \sigma_i(X_u)-\sigma_i(\bar{Y}_u)+\tilde{R}_1(\sigma_i)\Big\rangle dW_u^i \Big\|_{L^{\frac{p}{2}}(\Omega;\mathbb{R})}\\
\le\;& p\sum_{i=1}^{m} \Big(\int_0^t \Big\| \Big\langle X_s-\bar{Y}_s, \sigma_i(X_s)-\sigma_i(\bar{Y}_s)+\tilde{R}_1(\sigma_i)\Big\rangle \Big\|^2_{L^{\frac{p}{2}}(\Omega;\mathbb{R})} ds\Big)^{\frac{1}{2}}\\
\le\;& p\sum_{i=1}^{m} \Big(\int_0^t \|X_s-\bar{Y}_s\|^2_{L^p(\Omega;\mathbb{R}^d)} \cdot \Big\| \sigma_i(X_s)-\sigma_i(\bar{Y}_s)+\tilde{R}_1(\sigma_i)\Big\|^2_{L^p(\Omega;\mathbb{R}^d)} ds\Big)^{\frac{1}{2}}\\
\le\;& \sup_{0\le s\le t} \|X_s-\bar{Y}_s\|_{L^p(\Omega;\mathbb{R}^d)} \cdot p\sum_{i=1}^{m} \Big(\int_0^t \Big\| \sigma_i(X_s)-\sigma_i(\bar{Y}_s)+\tilde{R}_1(\sigma_i)\Big\|^2_{L^p(\Omega;\mathbb{R}^d)} ds\Big)^{\frac{1}{2}}\\
\le\;& \frac{1}{4}\sup_{0\le s\le t} \|X_s-\bar{Y}_s\|^2_{L^p(\Omega;\mathbb{R}^d)} + p^2 m\sum_{i=1}^{m} \int_0^t \Big\| \sigma_i(X_s)-\sigma_i(\bar{Y}_s)+\tilde{R}_1(\sigma_i)\Big\|^2_{L^p(\Omega;\mathbb{R}^d)} ds\\
\le\;& \frac{1}{4}\sup_{0\le s\le t} \|X_s-\bar{Y}_s\|^2_{L^p(\Omega;\mathbb{R}^d)} + 2p^2m^2K\int_0^t \|X_s-\bar{Y}_s\|^2_{L^p(\Omega;\mathbb{R}^d)} ds\\
&+2p^2m\sum_{i=1}^{m}\int_0^t \|\tilde{R}_1(\sigma_i)\|^2_{L^p(\Omega;\mathbb{R}^d)} ds.
\end{aligned}
\tag{3.33}
$$

At the same time, replacing f in (3.9) by μ and using the Cauchy-Schwarz inequality and elementary inequalities give

$$
\begin{aligned}
&2\Big\| \sup_{0\le s\le t} \int_0^s \Big\langle X_u-\bar{Y}_u, \mu(\bar{Y}_u)-\mu(Y_{n_u})\Big\rangle du \Big\|_{L^{\frac{p}{2}}(\Omega;\mathbb{R})}\\
=&2\Big\| \sup_{0\le s\le t} \int_0^s \Big\langle X_u-\bar{Y}_u, \mu'(Y_{n_u})\big(\sigma(Y_{n_u})\Delta W_u\big)+\tilde{R}_1(\mu)\Big\rangle du \Big\|_{L^{\frac{p}{2}}(\Omega;\mathbb{R})}\\
\le& J+2\Big\| \sup_{0\le s\le t} \int_0^s \Big\langle X_u-\bar{Y}_u, \tilde{R}_1(\mu)\Big\rangle du \Big\|_{L^{\frac{p}{2}}(\Omega;\mathbb{R})}\\
\le& J+\int_0^t \|X_s-\bar{Y}_s\|^2_{L^p(\Omega;\mathbb{R}^d)} ds+\int_0^t \|\tilde{R}_1(\mu)\|^2_{L^p(\Omega;\mathbb{R}^d)} ds,
\end{aligned}
\tag{3.34}
$$

where we denote

$$
J=2\Big\| \sup_{0\le s\le t} \int_0^s \Big\langle X_u-\bar{Y}_u, \mu'(Y_{n_u})\sigma(Y_{n_u})\Delta W_u\Big\rangle du \Big\|_{L^{\frac{p}{2}}(\Omega;\mathbb{R})}. \tag{3.35}
$$

Inserting (3.33) and (3.34) to (3.32) yields

$$
\begin{aligned}
&\frac{3}{4}\Big\| \sup_{0\le s\le t} \|X_s - \bar{Y}_s\| \Big\|^2_{L^p(\Omega;\mathbb{R})} \\
\le& 2(K+1+mK^2+p^2m^2K)\int_0^t \|X_s-\bar{Y}_s\|^2_{L^p(\Omega;\mathbb{R}^d)}ds + h^2\int_0^t \|\mu(Y_{n_s})\|^4_{L^{2p}(\Omega;\mathbb{R}^d)}ds \qquad (3.36)\\
&+2(1+p^2m)\sum_{i=1}^m \int_0^t \|\tilde{R}_1(\sigma_i)\|^2_{L^p(\Omega;\mathbb{R}^d)}ds + \int_0^t \|\tilde{R}_1(\mu)\|^2_{L^p(\Omega;\mathbb{R}^d)}ds + J.
\end{aligned}
$$

Therefore it remains to estimate J as (3.35). Using (3.3), (3.19) and (3.12) shows

$$
\begin{aligned}
X_u - \bar{Y}_u =& X_{t_{n_u}} - Y_{n_u} + \int_{t_{n_u}}^u \mu(X_r)dr - \int_{t_{n_u}}^u \tilde{\mu}(Y_{n_u})dr \\
&+\sum_{i=1}^m \int_{t_{n_u}}^u \Big[\sigma_i(X_r) - \sigma_i(Y_{n_r}) - \sum_{j=1}^m L^j\sigma_i(Y_{n_r})\Delta W_r^j\Big] dW_r^i \\
=& \int_{t_{n_u}}^u \big[\mu(X_r) - \mu(X_{t_{n_u}})\big]dr + \sum_{i=1}^m \int_{t_{n_u}}^u \big[\sigma_i(X_r) - \sigma_i(\bar{Y}_r)\big]dW_r^i + \sum_{i=1}^m \int_{t_{n_u}}^u \tilde{R}_1(\sigma_i)dW_r^i \\
&+ (u-t_{n_u})\,\mu(X_{t_{n_u}}) - (u-t_{n_u})\,\tilde{\mu}(Y_{n_u}) + X_{t_{n_u}} - Y_{n_u}.
\end{aligned}
$$

Thus

$$
\begin{aligned}
J \le& 2\Big\| \sup_{0\le s\le t} \int_0^s \Big\langle \int_{t_{n_u}}^u \big[\mu(X_r) - \mu(X_{t_{n_u}})\big]dr, \mu'(Y_{n_u})\sigma(Y_{n_u})\Delta W_u \Big\rangle du \Big\|_{L^{\frac{p}{2}}(\Omega;\mathbb{R})} \\
&+2\Big\| \sup_{0\le s\le t} \int_0^s \Big\langle \sum_{i=1}^m \int_{t_{n_u}}^u \big[\sigma_i(X_r) - \sigma_i(\bar{Y}_r)\big]dW_r^i, \mu'(Y_{n_u})\sigma(Y_{n_u})\Delta W_u \Big\rangle du \Big\|_{L^{\frac{p}{2}}(\Omega;\mathbb{R})} \\
&+2\Big\| \sup_{0\le s\le t} \int_0^s \Big\langle \sum_{i=1}^m \int_{t_{n_u}}^u \tilde{R}_1(\sigma_i)dW_r^i, \mu'(Y_{n_u})\sigma(Y_{n_u})\Delta W_u \Big\rangle du \Big\|_{L^{\frac{p}{2}}(\Omega;\mathbb{R})} \\
&+2\Big\| \sup_{0\le s\le t} \int_0^s \Big\langle \zeta_{n_u}, \mu'(Y_{n_u})\sigma(Y_{n_u})\Delta W_u \Big\rangle du \Big\|_{L^{\frac{p}{2}}(\Omega;\mathbb{R})} \\
&+2\Big\| \sup_{0\le s\le t} \int_0^s \Big\langle X_{t_{n_u}} - Y_{n_u}, \mu'(Y_{n_u})\sigma(Y_{n_u})\Delta W_u \Big\rangle du \Big\|_{L^{\frac{p}{2}}(\Omega;\mathbb{R})}, \\
:=& J_1 + J_2 + J_3 + J_4 + J_5, \qquad (3.37)
\end{aligned}
$$

where $\zeta_{n_u} \in \mathcal{F}_{t_{n_u}}$ is defined by

$$
\zeta_{n_u} = (u-t_{n_u})\,\mu(X_{t_{n_u}}) - (u-t_{n_u})\,\tilde{\mu}(Y_{n_u}). \qquad (3.38)
$$

To begin with, we establish the estimate

$$
\|\mu'(Y_{n_u})\sigma(Y_{n_u})\Delta W_u\|_{L^p(\Omega;\mathbb{R}^d)} \le C_{p,T}h^{\frac{1}{2}}, \qquad (3.39)
$$

which is frequently used later. Using Hölder's inequality, Lemma 3.3 and Lemma 2.7, we know that for $0\le k\le N-1, t_k \le s < t_{k+1}$

$$
\|\mu'(Y_k)\sigma(Y_k)\|_{L^{2p}(\Omega;\mathbb{R}^{d\times m})} \le \|\mu'(Y_k)\|_{L^{4p}(\Omega;\mathbb{R}^{d\times d})}\|\sigma(Y_k)\|_{L^{4p}(\Omega;\mathbb{R}^{d\times m})} < \infty \qquad (3.40)
$$

and

$$\|W_s - W_{t_k}\|_{L^{2p}(\Omega;\mathbb{R}^m)} = \|W_{s-t_k} - W_{t_0}\|_{L^{2p}(\Omega;\mathbb{R}^m)} \leq 2p\sqrt{m}h^{1/2}. \tag{3.41}$$

Combining (3.40) and (3.41) one can readily obtain (3.39) by Hölder's inequality. Concerning J_1, we use Hölder's inequality, (3.16) and (3.39) to arrive at

$$J_1 \leq 2\int_0^t \int_{t_{n_u}}^u \Big\|\mu(X_r) - \mu(X_{t_{n_u}})\Big\|_{L^p(\Omega;\mathbb{R}^d)} \cdot \Big\|\mu'(Y_{n_u})\sigma(Y_{n_u})\Delta W_u\Big\|_{L^p(\Omega;\mathbb{R}^d)} drdu \leq C_{p,T}h^2. \tag{3.42}$$

For J_2, using Hölder's inequality, (2.2), (2.26), elementary inequalities and (3.39) gives

$$\begin{aligned}
J_2 &\leq 2\int_0^t \Big\|\sum_{i=1}^m \int_{t_{n_u}}^u \big[\sigma_i(X_r) - \sigma_i(\bar{Y}_r)\big]dW_r^i\Big\|_{L^p(\Omega;\mathbb{R}^d)} \cdot \Big\|\mu'(Y_{n_u})\sigma(Y_{n_u})\Delta W_u\Big\|_{L^p(\Omega;\mathbb{R}^d)} du \\
&\leq \int_0^t \frac{1}{h}\Big\|\sum_{i=1}^m \int_{t_{n_u}}^u \big[\sigma_i(X_r) - \sigma_i(\bar{Y}_r)\big]dW_r^i\Big\|^2_{L^p(\Omega;\mathbb{R}^d)} du + \int_0^t h\Big\|\mu'(Y_{n_u})\sigma(Y_{n_u})\Delta W_u\Big\|^2_{L^p(\Omega;\mathbb{R}^d)} du \\
&\leq \frac{p^2}{h}\int_0^t \int_{t_{n_u}}^u \sum_{i=1}^m \Big\|\sigma_i(X_r) - \sigma_i(\bar{Y}_r)\Big\|^2_{L^p(\Omega;\mathbb{R}^d)} drdu + C_{p,T}h^2 \\
&\leq mp^2K^2\int_0^t \sup_{0\leq r\leq u}\big\|X_r - \bar{Y}_r\big\|^2_{L^p(\Omega;\mathbb{R}^d)} du + C_{p,T}h^2.
\end{aligned} \tag{3.43}$$

For J_3, similarly as above we obtain that

$$\begin{aligned}
J_3 \leq& 2\int_0^t \Big\|\sum_{i=1}^m \int_{t_{n_u}}^u \tilde{R}_1(\sigma_i)dW_r^i\Big\|_{L^p(\Omega;\mathbb{R}^d)} \cdot \Big\|\mu'(Y_{n_u})\sigma(Y_{n_u})\Delta W_u\Big\|_{L^p(\Omega;\mathbb{R}^d)} du \\
\leq& 2p\int_0^t \Big(\int_{t_{n_u}}^u \sum_{i=1}^m \Big\|\tilde{R}_1(\sigma_i)\Big\|^2_{L^p(\Omega;\mathbb{R}^d)} dr\Big)^{\frac{1}{2}} \cdot \Big\|\mu'(Y_{n_u})\sigma(Y_{n_u})\Delta W_u\Big\|_{L^p(\Omega;\mathbb{R}^d)} du \\
\leq& C_{p,T}h^2,
\end{aligned} \tag{3.44}$$

where (3.20) and (3.39) were also used. Now, it remains to estimate J_4 and J_5. We split J_4 into two terms as follows:

$$\begin{aligned}
J_4 \leq& 2\bigg\|\sup_{0\leq s\leq t}\Big|\sum_{k=0}^{n_s-1}\int_{t_k}^{t_{k+1}} \Big\langle \zeta_k, \mu'(Y_k)\sigma(Y_k)\Delta W_u\Big\rangle du\Big|\bigg\|_{L^{\frac{p}{2}}(\Omega;\mathbb{R})} \\
&+ 2\bigg\|\sup_{0\leq s\leq t}\Big|\int_{t_{n_s}}^s \Big\langle \zeta_{n_s}, \mu'(Y_{n_s})\sigma(Y_{n_s})\Delta W_u\Big\rangle du\Big|\bigg\|_{L^{\frac{p}{2}}(\Omega;\mathbb{R})} \\
:=& J_{41} + J_{42}.
\end{aligned} \tag{3.45}$$

Recall that $\zeta_k \in \mathcal{F}_{t_k}$ for $k = 0, 1, ..., N-1$. It can be readily verified that the discrete time process

$$\chi_n := \left\{\sum_{k=0}^{n-1}\int_{t_k}^{t_{k+1}} \Big\langle \zeta_k, \mu'(Y_k)\sigma(Y_k)\Delta W_u\Big\rangle du\right\}, \ \ n \in \{0, 1, ..., N\}$$

is an $\{\mathcal{F}_{t_n} : 0 \leq n \leq N\}$-martingale. With the aid of Doob's maximal inequality, Lemma 3.7 and Hölder's inequality we obtain that for $p \geq 4$

$$\begin{aligned} J_{41} &\leq \frac{2p}{p-2} \Big\| \sum_{k=0}^{n_t-1} \int_{t_k}^{t_{k+1}} \Big\langle \zeta_k, \mu'(Y_k)\sigma(Y_k)\Delta W_u \Big\rangle du \Big\|_{L^{\frac{p}{2}}(\Omega;\mathbb{R})} \\ &\leq \frac{2pc_{p/2}}{p-2} \Big(\sum_{k=0}^{n_t-1} \Big\| \int_{t_k}^{t_{k+1}} \Big\langle \zeta_k, \mu'(Y_k)\sigma(Y_k)\Delta W_u \Big\rangle du \Big\|^2_{L^{\frac{p}{2}}(\Omega;\mathbb{R})} \Big)^{1/2} \\ &\leq \frac{2pc_{p/2}}{p-2} \Big(\sum_{k=0}^{n_t-1} h \int_{t_k}^{t_{k+1}} \Big\| \Big\langle \zeta_k, \mu'(Y_k)\sigma(Y_k)\Delta W_u \Big\rangle \Big\|^2_{L^{\frac{p}{2}}(\Omega;\mathbb{R})} du \Big)^{1/2} \\ &\leq \frac{2pc_{p/2}}{p-2} \Big(\sum_{k=0}^{n_t-1} h \int_{t_k}^{t_{k+1}} \big\|\zeta_k\big\|^2_{L^p(\Omega;\mathbb{R}^d)} \cdot \big\|\mu'(Y_k)\sigma(Y_k)\Delta W_u\big\|^2_{L^p(\Omega;\mathbb{R}^d)} du \Big)^{1/2}, \end{aligned} \tag{3.46}$$

where by (3.38), Lemma 3.3 and Lemma 3.4

$$\big\|\zeta_k\big\|_{L^p(\Omega;\mathbb{R}^d)} \leq h\|\mu(X_{t_k})\|_{L^p(\Omega;\mathbb{R}^d)} + h\|\tilde{\mu}(Y_k)\|_{L^p(\Omega;\mathbb{R}^d)} \leq C_{p,T}h. \tag{3.47}$$

Hence, taking (3.39) and (3.47) into account, we derive from (3.46) that

$$J_{41} \leq C_{p,T}h^2. \tag{3.48}$$

For the second term J_{42}, Hölder's inequality, (3.47) and (3.39) give that for $p \geq 4$

$$\begin{aligned} \Big(\frac{J_{42}}{2}\Big)^{\frac{p}{2}} =& \mathbb{E}\Big(\sup_{0\leq s\leq t} \Big| \int_{t_{n_s}}^{s} \langle \zeta_{n_s}, \mu'(Y_{n_s})\sigma(Y_{n_s})\Delta W_u \rangle du \Big| \Big)^{\frac{p}{2}} \\ \leq& h^{\frac{p}{2}-1}\mathbb{E}\Big(\sup_{0\leq s\leq t} \int_{t_{n_s}}^{s} \Big| \langle \zeta_{n_s}, \mu'(Y_{n_s})\sigma(Y_{n_s})\Delta W_u \rangle \Big|^{\frac{p}{2}} du \Big) \\ \leq& h^{\frac{p}{2}-1}\mathbb{E}\Big(\sum_{k=0}^{n_t-1} \int_{t_k}^{t_{k+1}} \Big| \langle \zeta_k, \mu'(Y_k)\sigma(Y_k)\Delta W_u \rangle \Big|^{\frac{p}{2}} du + \int_{t_{n_t}}^{t} \Big| \langle \zeta_{n_s}, \mu'(Y_{n_s})\sigma(Y_{n_s})\Delta W_u \rangle \Big|^{\frac{p}{2}} du \Big) \\ =& h^{\frac{p}{2}-1} \int_0^t \mathbb{E}\Big| \langle \zeta_{n_u}, \mu'(Y_{n_u})\sigma(Y_{n_u})\Delta W_u \rangle \Big|^{\frac{p}{2}} du \\ \leq& h^{\frac{p}{2}-1} \int_0^t \|\zeta_{n_u}\|^{\frac{p}{2}}_{L^p(\Omega;\mathbb{R}^d)} \cdot \|\mu'(Y_{n_u})\sigma(Y_{n_u})\Delta W_u\|^{\frac{p}{2}}_{L^p(\Omega;\mathbb{R}^d)} du \\ \leq& C_{p,T}h^{\frac{5p}{4}-1}, \end{aligned} \tag{3.49}$$

which implies that for $p \geq 4$ and $h \in (0,1]$ there exists a suitable constant $C_{p,T}$ such that

$$J_{42} \leq 2C_{p,T}^{\frac{2}{p}} h^{\frac{5p-4}{2p}} \leq C_{p,T}h^2. \tag{3.50}$$

Gathering (3.48) and (3.50) we derive from (3.45) that for $p \geq 4$

$$J_4 \leq C_{p,T}h^2. \tag{3.51}$$

Similarly, we split J_5 as follows:

$$\begin{aligned} J_5 \leq & 2 \Big\| \sup_{0\leq s\leq t} \Big| \sum_{k=0}^{n_s-1} \int_{t_k}^{t_{k+1}} \Big\langle X_{t_k} - Y_k, \mu'(Y_k)\sigma(Y_k)\Delta W_u \Big\rangle du \Big| \Big\|_{L^{\frac{p}{2}}(\Omega;\mathbb{R})} \\ & + 2 \Big\| \sup_{0\leq s\leq t} \Big| \int_{t_{n_s}}^{s} \Big\langle X_{t_{n_s}} - Y_{n_s}, \mu'(Y_{n_s})\sigma(Y_{n_s})\Delta W_u \Big\rangle du \Big| \Big\|_{L^{\frac{p}{2}}(\Omega;\mathbb{R})} \\ := & J_{51} + J_{52}. \end{aligned} \tag{3.52}$$

With regard to J_{51}, following the same line as (3.46) yields

$$\begin{aligned} J_{51} \leq & \frac{2pc_{p/2}}{p-2} \Big(\sum_{k=0}^{n_t-1} h \int_{t_k}^{t_{k+1}} \|X_{t_k} - Y_k\|^2_{L^p(\Omega;\mathbb{R}^d)} \cdot \|\mu'(Y_k)\sigma(Y_k)\Delta W_u\|^2_{L^p(\Omega;\mathbb{R}^d)} du \Big)^{1/2} \\ \leq & \sup_{0\leq s\leq t} \|X_s - \bar{Y}_s\|_{L^p(\Omega;\mathbb{R}^d)} \cdot \frac{2pc_{p/2}}{p-2} \Big(\sum_{k=0}^{n_t-1} h \int_{t_k}^{t_{k+1}} \|\mu'(Y_k)\sigma(Y_k)\Delta W_u\|^2_{L^p(\Omega;\mathbb{R}^d)} du \Big)^{1/2} \\ \leq & \frac{1}{4} \sup_{0\leq s\leq t} \|X_s - \bar{Y}_s\|^2_{L^p(\Omega;\mathbb{R}^d)} + \frac{4p^2c^2_{p/2}}{(p-2)^2} \sum_{k=0}^{n_t-1} h \int_{t_k}^{t_{k+1}} \|\mu'(Y_k)\sigma(Y_k)\Delta W_u\|^2_{L^p(\Omega;\mathbb{R}^d)} du \\ \leq & \frac{1}{4} \sup_{0\leq s\leq t} \|X_s - \bar{Y}_s\|^2_{L^p(\Omega;\mathbb{R}^d)} + C_{p,T} h^2, \end{aligned} \tag{3.53}$$

where the fact $\bar{Y}_{t_k} = Y_k, k = 0, 1, ..., N-1$, elementary inequalities and (3.39) were used. For J_{52}, we follow the same way as (3.49) to obtain

$$\begin{aligned} \Big(\frac{J_{52}}{2}\Big)^{\frac{p}{2}} \leq & h^{\frac{p}{2}-1} \int_0^t \|X_{t_{n_u}} - Y_{n_u}\|^{\frac{p}{2}}_{L^p(\Omega;\mathbb{R}^d)} \cdot \|\mu'(Y_{n_u})\sigma(Y_{n_u})\Delta W_u\|^{\frac{p}{2}}_{L^p(\Omega;\mathbb{R}^d)} du \\ \leq & \sup_{0\leq s\leq t} \|X_s - \bar{Y}_s\|^{\frac{p}{2}}_{L^p(\Omega;\mathbb{R}^d)} \cdot C_{p,T} h^{\frac{3p}{4}-1}. \end{aligned} \tag{3.54}$$

Thus

$$J_{52} \leq 2 \sup_{0\leq s\leq t} \|X_s - \bar{Y}_s\|_{L^p(\Omega;\mathbb{R}^d)} \cdot C^{\frac{2}{p}}_{p,T}\, h^{\frac{3p-4}{2p}} \leq \frac{1}{4} \sup_{0\leq s\leq t} \|X_s - \bar{Y}_s\|^2_{L^p(\Omega;\mathbb{R}^d)} + 4C^{\frac{4}{p}}_{p,T}\, h^{\frac{3p-4}{p}}. \tag{3.55}$$

Note that $\frac{3p-4}{4} \geq 2$ for $p \geq 4$ and that $h \leq 1$. Plugging (3.53) and (3.55) into (3.52) yields

$$J_5 \leq \frac{1}{2} \sup_{0\leq s\leq t} \|X_s - \bar{Y}_s\|^2_{L^p(\Omega;\mathbb{R}^d)} + C_{p,T} h^2. \tag{3.56}$$

Inserting (3.42), (3.43), (3.44), (3.51) and (3.56) into (3.37) leads to

$$J \leq mp^2K^2 \int_0^t \sup_{0\leq r\leq u} \Big\| X_r - \bar{Y}_r \Big\|^2_{L^p(\Omega;\mathbb{R}^d)} du + \frac{1}{2} \sup_{0\leq s\leq t} \Big\| X_s - \bar{Y}_s \Big\|^2_{L^p(\Omega;\mathbb{R}^d)} + C_{p,T} h^2. \tag{3.57}$$

Hence, using estimates in Lemma 3.3 and Lemma 3.6 we derive from (3.36) that

$$\begin{aligned} \frac{3}{4} \Big\| \sup_{0\leq s\leq t} \|X_s - \bar{Y}_s\| \Big\|^2_{L^p(\Omega;\mathbb{R})} \leq & 2(K + 1 + mK^2 + p^2m^2K + \frac{1}{2}mp^2K^2) \int_0^t \sup_{0\leq u\leq s} \|X_u - \bar{Y}_u\|^2_{L^p(\Omega;\mathbb{R}^d)} ds \\ & + \frac{1}{2} \sup_{0\leq s\leq t} \Big\| X_s - \bar{Y}_s \Big\|^2_{L^p(\Omega;\mathbb{R}^d)} + C_{p,T} h^2. \end{aligned}$$

Thus $\left\| \sup_{0\le s\le t} \|X_s - \bar{Y}_s\| \right\|^2_{L^p(\Omega;\mathbb{R})}$ is finite by Lemma 3.4 and

$$\left\| \sup_{0\le s\le t} \|X_s - \bar{Y}_s\| \right\|^2_{L^p(\Omega;\mathbb{R})} \le C_{p,T} \int_0^t \left\| \sup_{0\le u\le s} \|X_u - \bar{Y}_u\| \right\|^2_{L^p(\Omega;\mathbb{R})} ds + C_{p,T} h^2. \tag{3.58}$$

The Gronwall inequality gives the desired result for $p \ge 4$. Using Hölder's inequality gives the assertion for $1 \le p < 4$ and the proof is complete. □

4 An illustrative example

In [9], the authors have demonstrated the computational efficiency of the tamed Euler scheme, compared to the implicit Euler method. In this section we compare computational efficiency of the tamed Milstein scheme and the tamed Euler scheme. To this end we choose a simple SDE (1.1)

$$dX_t = -X_t^5 dt + X_t dW_t, \quad X_0 = 1 \tag{4.1}$$

for $t \in [0,1]$. Figure 1 depicts the root mean-square errors (1.3) as a function of the stepsize h in log-log plot, where the expectation is approximated by the mean of 5000 independent realizations. As expected, the tamed Milstein scheme gives an error that decreases proportional to h, whereas the tamed Euler scheme gives errors that decrease proportional to $h^{\frac{1}{2}}$. To show the efficiency of the tamed Milstein method clearly, we present in Figure 2 the root mean-square errors of both methods as function of the runtime when $N \in \{2^{10}, ..., 2^{17}\}$ and the mean of 1000 independent paths are used to approximate the expectation in (1.3). Suppose that the strong approximation problem (1.3) of the SDE (4.1) should be solved with the precision $\varepsilon = 0.001$. From Figure 2, one can detect that $N = 2^{10}$ in the case of the tamed Milstein method (1.10) and that $N = 2^{16}$ in the case of the tamed Euler method (1.5) achieves the desired precision $\varepsilon = 0.001$ in (1.3). Moreover, the tamed Milstein scheme requires 8.1860 seconds while the tamed Euler scheme requires 147.9230 seconds to achieve the precision $\varepsilon = 0.001$ in (1.3). The tamed Milstein method is for the SDE (4.1) with commutative noise thus much faster than the tamed Euler method.

References

[1] A. Ambrosetti, and G. Prodi, A Primer of Nonlinear Analysis, Cambridge studies in advanced mathematics 34, CUP, 1993.

[2] G. Da Prato, and J. Zabczyk, Stochastic equations in infinite dimensions, vol. 44 of Encyclopedia of Mathematics and its Applications, Cambridge University Press, Cambridge, 1992.

[3] M. Giles, Multilevel Monte Carlo path simulation. Oper. Res. 56 (2008), pp. 607-617.

[4] D. Higham, X. Mao, and A. Stuart, Strong convergence of Euler-type methods for nonlinear stochastic diffrential equations. SIAM J. Numer. Anal. 40 (2002), pp. 1041-1063.



Figure 1: Root mean square approximation error versus stepsize h to approximate (4.1).



Figure 2: Root mean square approximation error versus runtime for $N \in \{2^{10}, ..., 2^{17}\}$.

[5] D. Higham, and P.E. Kloeden, Strong convergence rates for backward Euler on a class of nonlinear jump-diffusion problems, J. Comput. Appl. Math. 205 (2007), pp. 949-956.

[6] Y. Hu, Semi-implicit Euler-Maruyama scheme for stiff stochastic equations, in Stochastic Analysis and Related Topics V: The Silvri Workshop, Progr. Probab. 38 (1996), H. Koerezlioglu, ed., Birkhauser, Boston, pp. 183-202.

[7] M. Hutzenthaler, A. Jentzen, and P.E. Kloeden, Strong and weak divergence in finite time of Euler's method for SDEs with non-globally Lipschitz continuous coefficients, Proc. R. Soc. A 467 (2011), pp. 1563-1576.

[8] M. Hutzenthaler, and A. Jentzen, Convergence of the stochastic Euler scheme for locally Lipschitz Coefficients, Found. Comput. Math. 11 (2011), pp. 657-706.

[9] M. Hutzenthaler, A. Jentzen, and P.E. Kloeden, Strong convergence of an explicit numerical method for SDEs with non-globally Lipschitz continuous coefficients, arXiv:1010.3756v1, 2010.

[10] A. Jentzen, and M. Röckner, A Break of the Complexity of the Numerical Approximation of Nonlinear SPDEs with Multiplicative Noise, Arxiv preprint arXiv:1001.2751, 2010.

[11] I. Karatzas, S. Shreve, Brownian Motion and Stochatic Calculus. Springer, New York, 1991.

[12] P.E. Kloeden, and E. Platen, Numerical Solution of Stochastic Differential Equations, Springer, Berlin, 1992.

[13] P.E. Kloeden, E. Platen, and W. Wright, The approximation of multiple stochastic integrals. Stochastic Anal. Appl. 10 (1992), pp. 431-441.

[14] X. Mao, Stochastic Differential Equations and Applications, Horwood, New York, 1997.

[15] G. Maruyama, Continuous Markov processes and stochastic equations, Rend. Circolo. Math. Palermo 4 (1955), pp. 48-90.

[16] G. Milstein, Approximate integration of stochastic differential equations, Theor. Prob. Appl. 19 (1974), pp. 557-562.